arXiv:math/0602188v1 [math.PR] 9 Feb 2006

Isoperimetric-type inequalities for iterated Brownian motion in $\mathbb{R}^n$

Erkan Nane*
Department of Mathematics
Purdue University
West Lafayette, IN 47906
enane@math.purdue.edu

February 2, 2008

Abstract

We extend generalized isoperimetric-type inequalities to iterated Brownian motion over several domains in $\mathbb{R}^n$. *These kinds of inequalities imply in particular that for domains of finite volume, the exit distribution and moments of the first exit time for iterated Brownian motion are maximized with the ball* D^* *centered at the origin, which has the same volume as* D.

Mathematics Subject Classification (2000): 60J65, 60K99.
Key words: Iterated Brownian motion, Brownian-time Brownian motion, exit time, bounded domain, isoperimetric inequality.

---

[*] Supported in part by NSF Grant # 9700585-DMS

1 Introduction

Isoperimetric inequalities for Brownian motion and several other processes are well-known results. These kinds of inequalities give information about the geometry of the domain (see [2], [3], [9]-[11], [12]). Our aim in this paper is to extend these generalized isoperimetric inequalities to iterated Brownian motion (IBM) over several domains in $\mathbb{R}^n$.

To define the iterated Brownian motion Z_t started at $z \in \mathbb{R}$, let X_t^+, X_t^- and Y_t be three independent one-dimensional Brownian motions, all started at 0. Two-sided Brownian motion is defined by

$$X_t = \begin{cases} X_t^+, & t \geq 0 \\ X_{(-t)}^-, & t < 0. \end{cases}$$

Then the iterated Brownian motion started at $z \in \mathbb{R}$ is

$$Z_t = z + X(Y_t), \quad t \geq 0.$$

In $\mathbb{R}^n$, one requires $X^{\pm}$ to be independent $n-$dimensional Brownian motions. This is the version of the iterated Brownian motion due to Burdzy, see [4].

Properties of IBM and Brownian-time Brownian motion (BTBM), defined below, analogous to the properties of Brownian motion have been studied extensively by several authors (see [4, 5], [7, 8], [13, 14], [15] and the references therein). One of the main differences between these iterated processes and Brownian motion is that they are not Markov processes. However, these processes have connections with the parabolic operator $\frac{1}{8}\Delta^2 - \frac{\partial}{\partial t}$, as described in [1, 6].

We next give the background on isoperimetric inequalities that motivates us to study generalized isoperimetric inequalities for iterated Brownian motion.

Let $D \subset \mathbb{R}^n$ be a domain of finite volume, and denote by D^* the ball in $\mathbb{R}^n$ centered at the origin with the same volume as D. The class of quantities related to the Dirichlet Laplacian in D which are maximized or minimized by the corresponding quantities for D^* are often called generalized isoperimetric inequalities (C. Bandle [2]).

Probabilistically generalized isoperimetric inequalities read as

$$P_z[\tau_D > t] \;\leq\; P_0[\tau_{D^*} > t] \tag{1.1}$$

for all $z \in D$ and all $t > 0$, where τ_D is the first exit time of the Brownian motion from the domain D and P_z is the associated probability measure when this process starts at z. These kinds of Isoperimetric inequalities imply eigenvalue inequalities for Dirichlet Laplacian, Integrals of Green functions of the Dirichlet Laplacian and the moments of the exit time of Brownian motion (see [9] - [11]). These inequalities serve as motivation for our results in this paper.

Isoperimetric inequalities have been extended in many directions including Brownian motion over convex domains in $\mathbb{R}^n$ where, one fixes the inradius R_D of D, instead of fixing the volume. The inradius is the supremum of the radius of all the balls contained in D. Indeed, a special case of the theorem proved by R. Bañuelos, R. Latała, and P. Meǹdez in [3] is the following; suppose $D \subset \mathbb{R}^2$ is a convex domain of finite inradius R_D. Then

$$P_z[\tau_D > t] \;\leq\; P_0[\tau_{S(D)} > t] \;=\; P_0[\tau_{I(D)} > t] \tag{1.2}$$

for all $z \in D$ and all $t > 0$ where, $I(D) = (-R_D, R_D)$ and $S(D) = \mathbb{R} \times I(D)$.

More recently, P. Meǹdez [12] proved a generalization of inequality (1.2) in $\mathbb{R}^n$. Suppose $D \subset \mathbb{R}^n$ is a convex domain of finite inradius R_D. Then

$$P_z[\tau_D > t] \;\leq\; P_0[\tau_{S(D)} > t] \;=\; P_0[\tau_{I(D)} > t] \tag{1.3}$$

for all $z \in D$ and all $t > 0$ where, $I(D) = (-R_D, R_D)$ and $S(D) = \mathbb{R}^{n-1} \times I(D)$.

P. Meǹdez proved in [12, Theorem 5.2] a sharper version of inequality (1.2) in $\mathbb{R}^2$. Suppose $D \subset \mathbb{R}^2$ is a convex domain of finite inradius R_D and finite diameter d_D. Then

$$P_z[\tau_D > t] \;\leq\; P_0[\tau_{C(D)} > t] \tag{1.4}$$

for all $z \in D$ and all $t > 0$ where, $C(D) = [\mathbb{R} \times I(D)] \cap B(0, d_D \;-\; R_D)$.

In analogy with ordinary Brownian motion and diffusions, if $\tau_D(Z)$ is the first exit time of IBM from domain D, started at $z \in D$,

$$(i.e.\;\; \tau_D(Z) = \inf\{t \geq 0 : \; Z_t \notin D\}).$$

Then $P_z[\tau_D(Z) > t]$ provides a measure of the lifetime of the process in D.

Let $D \subset \mathbb{R}^n$ be an open set and let D_* be any of the sets D^*, $S(D)$, $I(D)$ or $C(D)$. $D_* = D^*$ the open ball of the same volume as D when D is of finite volume. $D_* = S(D) = \mathbb{R}^{n-1} \times I(D)$ where $I(D) = (-R_D, R_D)$ when D is a convex domain of finite inradius R_D. $D_* = C(D) = [\mathbb{R} \times I(D)] \cap B(0, d_D - R_D)$ when $D \subset \mathbb{R}^2$ is a convex set of finite inradius R_D and finite diameter d_D.

The following is the main result of this paper. It is an extension of generalized isoperimetric-type inequalities to iterated Brownian motion over several domains.

Theorem 1.1. *Let $D \subset \mathbb{R}^n$ be an open set, of finite volume or a convex set of finite inradius or a convex set of finite inradius with finite diameter, then*

$$P_z[\tau_D(Z) > t] \;\leq\; P_0[\tau_{D_*}(Z) > t] \tag{1.5}$$

Let ϕ be an increasing function. Multiplying the inequality (1.5) by the derivative of ϕ and integrating the resulting inequality in time from zero to infinity, we obtain the following corollary.

Corollary 1.1. *Let ϕ be a nonnegative increasing function, then*

$$E_z(\phi(\tau_D(Z))) \leq E_0(\phi(\tau_{D_*}(Z))) \tag{1.6}$$

In particular Corollary 1.1 implies that for $p \geq 1$,

$$E_z((\tau_D(Z)))^p \leq E_0((\tau_{D_*}(Z))^p) \tag{1.7}$$

Theorem 1.1, and its corollaries imply that the distribution functions, and the moments of the exit times are maximized with D^* when D is an open set of finite volume, with $S(D)$ when D is a convex domain of finite inradius and with $C(D)$ when D is a convex domain of finite inradius and finite diameter.

We also obtain a version of the above Theorem 1.1 for another closely related process, the so called Brownian-time Brownian motion (BTBM). To define this, let X_t and Y_t be two independent one-dimensional Brownian motions, all started at 0. BTBM is defined to be $Z^1_t = x + X(|Y_t|)$. Properties of this process and its connections to PDE's have been studied in [1] and [8].

In analogy with ordinary Brownian motion and diffusions, if $\tau_D(Z^1)$ is the first exit time of BTBM from domain D, started at $z \in D$, $P_z[\tau_D(Z^1) > t]$ provides a measure of the lifetime of the process in D. Then we have

Theorem 1.2. *Let $D \subset \mathbb{R}^n$ be an open set, of finite volume or a convex set of finite inradius or a convex set of finite inradius with finite diameter, then*

$$P_z[\tau_D(Z^1) > t] \;\leq\; P_0[\tau_{D_*}(Z^1) > t] \tag{1.8}$$

Let ϕ be an increasing function. Multiplying the inequality (1.8) by the derivative of ϕ and integrating the resulting inequality in time from zero to infinity, we obtain the following corollary.

Corollary 1.2. *Let ϕ be a nonnegative increasing function, then*

$$E_z(\phi(\tau_D(Z^1))) \leq E_0(\phi(\tau_{D_*}(Z^1))) \tag{1.9}$$

In particular Corollary 1.2 implies that for $p \geq 1$,

$$E_z((\tau_D(Z^1)))^p \leq E_0((\tau_{D_*}(Z^1))^p) \tag{1.10}$$

Theorem 1.2, and its corollaries imply that the distribution functions, and the moments of the exit times are maximized with D^* when D is an open set of finite volume, with $S(D)$ when D is a convex domain of finite inradius and with $C(D)$ when D is a convex domain of finite inradius and finite diameter.

We use integration by parts and the double integral representation of the exit distribution of the first exit time of IBM from domains in $\mathbb{R}^n$ to prove Theorems 1.1 and 1.2 in the next section.

2 Proofs of Main results

If $D \subset \mathbb{R}^n$ is an open set, write

$$\tau_D^{\pm}(z) = \inf\{t \geq 0 : \quad X_t^{\pm} + z \notin D\},$$

and if $I \subset \mathbb{R}$ is an open interval, write

$$\eta_I = \eta(I) = \inf\{t \geq 0 : \quad Y_t \notin I\}.$$

So we have in the notation of the introduction

$$P[\tau_D(z) > t] = P_z[\tau_D > t].$$

Recall that $\tau_D(Z)$ stands for the first exit time of iterated Brownian motion from D. As in DeBlassie [6, §3.], we have by the continuity of the paths for $Z_t = z + X(Y_t)$, if f is the probability density of $\tau_D^{\pm}(z)$

$$P_z[\tau_D(Z) > t] = \int_0^\infty \int_0^\infty P_0[\eta_{(-u,v)} > t] f(u) f(v) dv du. \tag{2.1}$$

To prove isoperimetric-type inequalities we need the following Lemmas.

Lemma 2.1. *For any* $u, v \in (0, \infty)$

$$\left(\frac{\partial}{\partial v} P_0[\eta_{(-u,v)} > t]\right) \geq 0,$$

and

$$\left(\frac{\partial}{\partial u} P_0[\eta_{(-u,v)} > t]\right) \geq 0$$

Proof. Fix $u \in (0, \infty)$, let $0 < v_1 \leq v_2 < \infty$. We have $(-u, v_1) \subset (-u, v_2)$, hence $\eta_{(-u,v_1)} \leq \eta_{(-u,v_2)}$. So, for any $t > 0$, $P_0[\eta_{(-u,v_1)} > t] \leq P_0[\eta_{(-u,v_2)} > t]$. Hence the first inequality holds. The last inequality is proved similarly.

□

Lemma 2.2 (Integration by parts lemma). *Let* D *be a domain such that* $\lim_{t\to\infty} P_z[\tau_D > t] = 0$. *Then*

$P_z[\tau_D(Z) > t]$ *is equal to*

$$= \int_0^\infty \int_0^\infty P_0[\eta_{(-u,v)} > t] f(u) f(v) dv du. \tag{2.2}$$

$$= \int_0^\infty \int_0^\infty \left(\frac{\partial}{\partial v} P_0[\eta_{(-u,v)} > t]\right) P[\tau_D(z) > v] f(u) dv du \tag{2.3}$$

$$= \int_0^\infty \int_0^\infty \left(\frac{\partial}{\partial u} P_0[\eta_{(-u,v)} > t]\right) P[\tau_D(z) > u] f(v) du dv. \tag{2.4}$$

Proof. Equation (2.3) follows by integration by parts and observing

$$-\frac{\partial}{\partial v} P[\tau_D(z) > v] = f(v)$$

and

$$\lim_{u\to 0} P_0[\eta_{(-u,v)} > t] P[\tau_D(z) > u] = \lim_{u\to\infty} P_0[\eta_{(-u,v)} > t] P[\tau_D(z) > u] = 0.$$

Equation (2.4) follows by Fubini-Tonelli theorem in equation (2.2) and by similar arguments as in the proof of equation (2.3). □

If we use integration by parts one more time, we obtain

$P_z[\tau_D(Z) > t]$

$$= \int_0^\infty \int_0^\infty \left(\frac{\partial}{\partial u}\frac{\partial}{\partial v} P_0[\eta_{(-u,v)} > t]\right) P[\tau_D(z) > u] P[\tau_D(z) > v] dv du. \tag{2.5}$$

We now can try to use the isoperimetric-type inequalities for Brownian motion to prove Theorem 1.1. This is valid if one can show that

$$\frac{\partial}{\partial u}\frac{\partial}{\partial v}P_0[\eta_{(-u,v)} > t] \geq 0$$

for all $u, v, t \in (0, \infty)$. But this is not easy to show (maybe not true), since this is given by an infinite series given in [13]:

$$\begin{aligned}
&\frac{\partial}{\partial u}\frac{\partial}{\partial v}P_0[\eta_{(-u,v)} > t] \\
&= 4\sum_{n=0}^{\infty} \exp\left(-\frac{(2n+1)^2\pi^2}{2(u+v)^2}t\right)\Bigg\{\Bigg[\sin\ \frac{(2n+1)\pi u}{u+v}\ \left(\frac{1}{(u+v)^4}\right) \\
&\qquad \times \left(\frac{\pi^3(2n+1)^3t^2}{(u+v)^2} - 3\pi(2n+1)t + (2n+1)\pi uv\right)\Bigg] \\
&+ \Bigg[\cos\ \frac{(2n+1)\pi u}{u+v}\ \left(\frac{1}{(u+v)^3}\right)\left(\frac{(2n+1)^2\pi^2t(v-u)}{(u+v)^2} + u - v\right)\Bigg]\Bigg\}.
\end{aligned}$$

So we follow another path to prove isoperimetric inequalities.

Proof of Theorem 1.1. The idea of the proof is using integration by parts and the corresponding generalized isoperimetric inequalities for Brownian motion. Let f^* denote the probability density of τ_{D_*}. By equation (2.1) and integration by parts $P_z[\tau_D(Z) > t]$ equals

$$\begin{aligned}
&\int_0^\infty\int_0^\infty \left(\frac{\partial}{\partial v}P_0[\eta_{(-u,v)} > t]\right)P[\tau_D(z) > v]f(u)dvdu && (2.6)\\
\leq\ &\int_0^\infty\int_0^\infty \left(\frac{\partial}{\partial v}P_0[\eta_{(-u,v)} > t]\right)P[\tau_{D_*}(0) > v]f(u)dvdu && (2.7)\\
=\ &\int_0^\infty\int_0^\infty P_0[\eta_{(-u,v)} > t]f(u)f^*(v)dvdu && (2.8)\\
=\ &\int_0^\infty\int_0^\infty \left(\frac{\partial}{\partial u}P_0[\eta_{(-u,v)} > t]\right)P[\tau_D(z) > u]f^*(v)dudv && (2.9)\\
\leq\ &\int_0^\infty\int_0^\infty \left(\frac{\partial}{\partial u}P_0[\eta_{(-u,v)} > t]\right)P[\tau_{D_*}(0) > v]f^*(u)dvdu && (2.10)\\
=\ &P_0[\tau_{D_*}(Z) > t] && (2.11)
\end{aligned}$$

Equations (2.7) and (2.10) follow from the corresponding isoperimetric inequalities for Brownian motion and Lemma 2.1. Equations (2.6), (2.8) and

(2.11) follow by integration by parts, Lemma 2.2. Equation (2.9) follows from Fubini-Tonelli theorem and integration by parts as in Lemma 2.2.

□

We next give the proof of Theorem 1.2. Let X_t and Y_t be two independent one-dimensional Brownian motions, all started at 0. BTBM is defined to be $Z_t^1 = x + X(|Y_t|)$. In $\mathbb{R}^n$, we require X to be independent one dimensional iterated Brownian motions. Let $\tau_D(Z^1)$ stand for the first exit time of BTBM from D. We have by the continuity of paths

$$P_z[\tau_D(Z^1) > t] = P[\eta(-\tau_D(z), \tau_D(z)) > t]. \tag{2.12}$$

To prove isoperimetric-type inequalities we need the following Lemma.

Lemma 2.3. *$\frac{\partial}{\partial u} P_0[\eta_{(-u,u)} > t] \geq 0$ for all $u > 0$.*

Proof. Since $P_0[\eta_{(-u,u)} > t] \leq P_0[\eta_{(-v,v)} > t]$ for $u \leq v$, the derivative is positive.

□

Proof of Theorem 1.2. From equation (2.12)

$$\begin{aligned} P_z[\tau_D(Z^1) > t] &= P[\eta(-\tau_D(z), \tau_D(z)) > t] \\ &= \int_0^\infty P_0[\eta_{(-u,u)} > t] f(u) du \qquad (2.13) \end{aligned}$$

By integration by parts, equation (2.13) equals

$$\int_0^\infty \frac{\partial}{\partial u} P_0(\eta_{(-u,u)} > t) P[\tau_D(z) > u] du$$

Now from Lemma 2.3, $\frac{\partial}{\partial u} P_0(\eta_{(-u,u)} > t) \geq 0$ for all $u > 0$, which gives the desired conclusion from the corresponding isoperimetric inequalities for the first exit time of the Brownian motion X from D_*.

□

We have a general result which follows from Theorems 1.2 and 1.1.

Corollary 2.1. *Let ξ and T be positive random variables such that*

$$\lim_{t\to\infty} P[\xi > t] = 0 = \lim_{t\to\infty} P[T > t]$$

and for each $t > 0$

$$P[\xi > t] \leq P[T > t].$$

Let ξ_1, ξ_2 *be independent copies of* ξ, T_1, T_2 *independent copies of* T *and let* Y *be a one dimensional Brownian motion independent of* ξ *and* T. *Then for each* $t > 0$,

$$P[\eta(-\xi_1, \xi_2) > t] \leq P[\eta(-T_1, T_2) > t]$$

and

$$P[\eta(-\xi, \xi) > t] \leq P[\eta(-T, T) > t].$$

Acknowledgments. I would like to thank Professor Rodrigo Bañuelos, my academic advisor, for suggesting this problem to me and for his guidance on this paper.

References

[1] H. Allouba and W. Zheng, *Brownian-time processes: The pde connection and the half-derivative generator,* Ann. Prob. 29 (2001), no. 2, 1780-1795.

[2] C. Bandle, *Isoperimetric Inequalities and Applications,* Monog. Stud. Math. 7, Pitnam, Boston, 1980.

[3] R. Bañuelos, R. Latała and P. J. Meǹdez-Hernaǹdez, *Brascamp-Lieb-Luttinger-type inequality and applications to symmetric stable processes,* Proc. Amer. Math. Soc. 129 (2001), 2997-3008.

[4] K. Burdzy, *Some path properties of iterated Brownian motion,* In Seminar on Stochastic Processes (E. Çinlar, K.L. Chung and M.J. Sharpe, eds.), Birkhäuser, Boston, (1993), 67-87.

[5] K. Burdzy, *Variation of iterated Brownian motion,* In Workshops and Conference on Measure-valued Processes, Stochastic Partial Differential Equations and Interacting Particle Systems (D.A. Dawson, ed.) Amer. Math. Soc. Providence, RI, (1994),35-53.

[6] R. D. DeBlassie, *Iterated Brownian motion in an open set,* Ann. Appl. Prob. 14 (2004), no. 3, 1529-1558.

[7] D. Khoshnevisan and T.M. Lewis, *Stochastic calculus for Brownian motion in a Brownian fracture,* Ann. Applied Probabl. 9 (1999), no. 3, 629-667.

[8] D. Khoshnevisan and T.M. Lewis, *Chung's law of the iterated logarithm for iterated Brownian motion,* Ann. Inst. H. Poincaré Probab. Statist. 32 (1996), no. 3, 349-359.

[9] J. M. Luttinger, *Generalized isoperimetric inequalities,* J. Math. Phys. **14** (1973), 586 - 593.

[10] J. M. Luttinger, *Generalized isoperimetric inequalities II,* J. Math. Phys. 14 (1973), 1444-1447.

[11] J. M. Luttinger, *Generalized isoperimetric inequalities III,* J. Math. Phys. 14 (1973), 1448-1450.

[12] P. J. Meǹdez-Hernaǹdez, *Brascamp-Lieb-Luttinger inequalities for convex domains of finite inradius,* Duke Math. J. 113 (2002), 93-131.

[13] E. Nane, *Iterated Brownian motion in parabola-shaped domains,* To appear in Potential Analysis.

[14] E. Nane, *Iterated Brownian motion in bounded domains in $\mathbb{R}^n$,* To appear in Stochastic Processes and Their Applications

[15] Y. Xiao, *Local times and related properties of multidimensional iterated Brownian motion,* J. Theoret. Probab. 11 (1998), no. 2, 383-408.